

Benchmarking Trisaminocyclopropeniums as Mediators for Anodic Oxidation Reactions

Sabrina N. Carneiro, Joshua D. Laffoon, Long Luo,* and Melanie S. Sanford*

Cite This: <https://doi.org/10.1021/acs.joc.4c00422>


Read Online

ACCESS |


Metrics & More

Article Recommendations

Supporting Information



Benchmark electrochemical properties,
performance as electron-transfer mediators



ABSTRACT: This report benchmarks a tris(amino)cyclopropenium (TAC) salt as an electron-transfer mediator for anodic oxidation reactions in comparison to two known mediators: a triarylamine and a triarylimidazole derivative. The three mediators have redox potentials, diffusion coefficients, and heterogeneous electron transfer rates similar to those of glassy carbon electrodes in acetonitrile/KPF₆. However, they differ significantly in their performance in two electro-organic reactions: anodic fluorination of a dithiane and anodic oxidation of 4-methoxybenzyl alcohol. These differences are rationalized based on variable stability in the presence of reaction components (e.g., NEt₃·3HF, lutidine, and Cs₂CO₃) as well as very different rates of electron transfer with the organic substrate. Overall, this work highlights the advantages and disadvantages of each mediator and provides a foundation for expanding the applications of TACs in electro-organic synthesis moving forward.

INTRODUCTION

Electrochemistry has emerged as a powerful modality in synthetic organic chemistry.¹ As such, there is increasing demand for tunable electron-transfer mediators that can effectively function as soluble electrodes during electro-organic redox reactions.² Mediated electrochemical transformations are highly desirable, as they often proceed at lower cell potentials, with higher yields and selectivities, and with less electrode passivation than the corresponding direct electrolysis pathways.² Several characteristics are required for an effective electron-transfer mediator, including (i) fast and reversible electron transfer with both solid electrodes and organic substrates, (ii) stability of all redox states to electro-organic reaction conditions, (iii) readily tunable redox potentials, and (iv) high solubility in all oxidation states. Oxidative mediators that meet most or all of these criteria while exhibiting relatively high redox potentials ($\geq +0.7$ V versus Ag/Ag⁺) remain rare. To date, the most widely used examples are polyaromatic molecules such as triarylamine³ or triarylimidazole⁴ derivatives (for example, A and B in Figure 1a). While these have been successfully employed as electron-transfer mediators for a variety of anodic oxidation reactions,² they can be challenging to synthesize/derivatize in a modular fashion, suffer from




Figure 1. (a) Known high-potential oxidative mediators for electro-organic synthesis. (b) This work: Tris(amino)cyclopropeniums as mediators for electro-organic synthesis.

Received: February 15, 2024

Revised: March 19, 2024

Accepted: March 26, 2024

modest solubility due to their propensity for aggregation and pi-stacking, and have variable stability under some electro-organic reaction conditions. Overall, there is a need for new high redox potential mediators for anodic oxidations that are readily synthetically tunable, are soluble in common organic media, and have complementary stability.

The Sanford group has recently developed cyclopropenium salts such as trisaminocyclopropenium (TAC) **C** (Figure 1b) as electrochemical energy storage materials in nonaqueous redox flow batteries (RFBs).⁵ We have disclosed a series of cyclopropenium derivatives that undergo oxidation at potentials ranging from +0.7 to 1.2 V versus Fc/Fc⁺ in acetonitrile (MeCN)/KPF₆.⁶ Many of the properties required for efficient energy storage (reversible redox events at high potentials accompanied by high stability and solubility) mirror those necessary for redox mediators.⁷ However, there are two key challenges for transitioning from energy storage materials to mediators in electro-organic synthesis. First, electrochemical cycling in RFBs occurs under highly controlled conditions: under an inert atmosphere with a rigorously purified and dried solvent and a supporting electrolyte. In contrast, electro-organic reactions are often conducted on the benchtop and contain a mixture of organic reactants and additives (e.g., acids, bases, water, and supporting salts). Second, in RFBs, electron transfer reactions occur exclusively at the solution–electrode interface. In contrast, effective redox mediators must undergo fast electron transfer with both the solid electrode and the relevant organic substrate.

In this report, we evaluate and compare cyclopropenium **C** (a high-potential, soluble TAC derivative)⁸ with triarylamine **A** and triarylimidazole **B** as redox mediators in organic synthesis. **A–C** have similar redox potentials ($E_{1/2} \approx 0.8$ V vs Ag/AgBF₄) and should thus be effective mediators for comparable classes of organic substrates. We first compare their electrochemical properties ($E_{1/2}$, diffusion, and electron transfer rates) under standard conditions and evaluate their compatibility with acidic and basic additives. Next, we assess their performance as mediators in two electro-organic reactions: the anodic fluorination of a dithiane⁹ and the anodic oxidation of 4-methoxybenzyl alcohol.^{4a} Overall, this work reveals advantages and disadvantages for each mediator and provides a foundation for expanding the applications of TACs in electro-organic synthesis moving forward.

RESULTS AND DISCUSSION


We initiated these studies by using cyclic voltammetry (CV) to compare the redox potentials ($E_{1/2}$), diffusion rates (D), and heterogeneous electron transfer rates (k_{het}) of **A–C** (Table 1 and Figures S1–S4). CVs were conducted with **A–C** (2 mM) in 0.5 M KPF₆ in MeCN using a glassy carbon working electrode, a platinum wire counter electrode, and a Ag/AgBF₄ reference electrode. KPF₆ was chosen as the supporting

electrolyte because it has been used extensively with TACs in flow batteries.⁵ Under these conditions, the $E_{1/2}$ values of **A–C** are within 140 mV of one another, ranging from 0.77 V (**A**) to 0.88 V (**B**) to 0.91 V (**C**) versus Ag/AgBF₄. At 100 mV/s, we observe reversible redox peaks for **A** and **C** ($|i_{\text{pa}}/i_{\text{pc}}| \approx 1$), while **B** displays quasi-reversibility that is exacerbated at slower scan rates ($|i_{\text{pa}}/i_{\text{pc}}| = 2.23$ at 10 mV/s). This suggests that the oxidized species **A**⁺ and **C**²⁺ are significantly longer-lived intermediates than **B**⁺.¹⁰

The rates of heterogeneous electron transfer for **A–C** at glassy carbon are similar to one another ($\sim 2 \times 10^{-2}$ cm/s), as determined using the Nicholson method.¹⁰ Notably, the values for **A** and **B** are similar to those reported by Little in 0.1 M LiClO₄ in MeCN, 1.38×10^{-2} and 1.53×10^{-2} cm/s, respectively.^{4c} The diffusion coefficients (D) for **A–C** were experimentally determined by applying the Randles–Sevcik eq (Figure S4).¹¹ **A** and **C** have diffusion coefficients of 1.4×10^{-5} cm²/s, while that for **B** is about 2-fold slower (6.58×10^{-6} cm²/s). Again, the values for **A** and **B** are similar to those determined by Little in 0.1 M LiClO₄ in MeCN.^{4c}

We next sought to benchmark **A–C** as mediators in two different electro-oxidative reactions: (a) the anodic fluorination of dithiane **1** with NEt₃·3HF to form **2** and (b) the anodic oxidation of 4-methoxybenzyl alcohol **3** to generate aldehyde **4** (Scheme 1). These transformations were selected for several

Scheme 1. Electrochemical Oxidation Reactions Used to Benchmark Mediators A–C



reasons. First, substrates **1** and **3** have peak anodic potentials (E_{pa}) of 1.00 and 1.23 V vs Ag/AgBF₄, respectively (Figure S1), which are well matched for mediation by **A–C**. Second, anodic fluorinations analogous to (a) have been studied with **A**,³ while benzyl alcohol oxidations analogous to (b) have been investigated with both **A** and **B**.² As such, these transformations offer an opportunity to compare the performance of **C** to that of the known mediators. Third, the conditions for these reactions are quite different, with (a) performed in mildly acidic (NEt₃·3HF) and (b) conducted in basic (lutidine) media. We hypothesized that these conditions might thus highlight complementary compatibility/performance between **A–C**.³

To test this latter point, we preliminarily examined the stability of **A–C** toward CV cycling in the absence and presence of NEt₃·3HF and lutidine as additives. CVs were conducted with **A–C** (2 mM) and the additive (10 mM) in 0.5 M KPF₆ in MeCN under ambient conditions.¹² In each case, 10 sequential CVs of the mediator were acquired at a rate of 100 mV/s. We then determined the % change in anodic peak current from cycle 2 to cycle 10 (% decrease of i_{pa}) to estimate stability toward each additive.¹³ The CVs from these experiments are listed in Figure S7, and the data are summarized in Table 2. Overall, the % decrease of i_{pa} varied

Table 1. Electrochemical Properties of A–C

	A	B	C
$E_{1/2}$ (V)	0.77	0.89	0.91
$ i_{\text{pa}}/i_{\text{pc}} $ (10 mV/s)	1.00	2.23	1.14
$ i_{\text{pa}}/i_{\text{pc}} $ (100 mV/s)	1.08	1.43	1.17
D [cm ² /s]	1.42×10^{-5}	6.48×10^{-6}	1.46×10^{-5}
k_{het} [cm/s]	2.2×10^{-2}	1.88×10^{-2}	1.76×10^{-2}
$k_{\text{ET-1}}$ [M ⁻¹ s ⁻¹]	22	6.3	140

Table 2. Stability of A–C towards $\text{NEt}_3\cdot 3\text{HF}$ and Lutidine, as Assessed by CV^a

entry	mediator	additive	% decrease of i_{pa}
1	A	none	2
2	A	$\text{NEt}_3\cdot 3\text{HF}$	4
3	A	lutidine	2
4	B	none	7
5	B	$\text{NEt}_3\cdot 3\text{HF}$	>19 ¹³
6	B	lutidine	8
7	C	none	9
8	C	$\text{NEt}_3\cdot 3\text{HF}$	6
9	C	lutidine	14

^aCV conditions: mediator (2 mM) and additive (10 mM) in 0.5 M KPF₆ in MeCN with a glassy carbon working electrode, a platinum wire counter electrode, and a Ag/AgBF₄ reference electrode. CVs acquired at 100 mV/s over 10 consecutive scans. %decrease in i_{pa} between cycles 2 and 10 is reported for each.

from 2 to >19%,¹³ and A–C exhibited complementary stability toward these additives. A and B showed the largest decrease in i_{pa} with $\text{NEt}_3\cdot 3\text{HF}$ (4% and >19%, respectively), while C showed the largest decrease with lutidine (14%).

The first step of the anodic fluorination of dithiane **1** involves electron transfer between the mediator and **1** to form the radical cation **1**^{•+} (Figure 2a). We sought to compare the rate constants of this electron transfer reaction ($k_{\text{ET-1}}$) for each of the mediators. Rate constants were obtained via foot-of-the-wave analysis¹⁴ of the CVs of A–C (2 mM in 0.5 M KPF₆ in MeCN) in the presence of 40 mM of dithiane **1** at scan rates ranging from 5 to 500 mV/s. During the foot-of-the-wave analysis, the CV data were postprocessed to generate the foot-of-the-wave analysis plots exemplified in Figure 2b for C. The y-axis is the ratio between the electrochemical current for C in the presence of substrate **1** (i) and the peak current for C in the absence of **1** (i_p^0) at the same scan rate. The x-axis is a function of the electrode potential (E): $\frac{1}{1 + \exp\left[\frac{F}{RT}(E - E^0)\right]}$, where

F is the Faraday constant (=96,480 C/mol), R is the gas constant, T is 298 K, and E^0 is the standard reduction potential of the mediator. The slope of the linear fit of the initial stage of the curve (i.e., foot-of-the-wave, highlighted in thick lines in Figure 2b) provides the rate constant ($k_{\text{ET-1}}$) based on eq 1, where C^0 is the concentration of substrate **1**, and ν is the scan rate.

$$\frac{i}{i_p^0} = \left(2.24 \sqrt{\frac{RT}{F\nu}} 2k_{\text{ET-1}}C^0\right) \frac{1}{1 + \exp\left[\frac{F}{RT}(E - E^0)\right]} \quad (1)$$

As shown in Figure 2c, the slopes at different scan rates obtained by linear fitting were proportional to $\nu^{-0.5}$, in excellent agreement with eq 1. The $k_{\text{ET-1}}$ values were calculated from the slopes of the dotted fit lines in Figure 2c. Large differences are observed between the three mediators, with $k_{\text{ET-1}}$ varying from 6.3 (B) to 22 (A) to 140 (C) $\text{M}^{-1} \text{s}^{-1}$. Notably, these values do not directly track with differences in redox potential,^{4c} suggesting that a more complex set of factors impacts $k_{\text{ET-1}}$. Overall, the large value of $k_{\text{ET-1}}$ with C is promising for the mediated electrolysis of dithiane **1** on the preparative scale.

The preparative scale A–C-catalyzed electrolysis reactions of **1** with $\text{NEt}_3\cdot 3\text{HF}$ were next evaluated in a divided cell equipped with Pt foil electrodes using 0.7 M $\text{NEt}_3\cdot 3\text{HF}$ as both the electrolyte and the fluorinating reagent in MeCN. Note






Figure 2. (a) Electron transfer between the oxidized mediator (mediator^{•+}) and substrate **1**. (b) Foot-of-the-wave analysis for mediator C. (c) Fitting to obtain $k_{\text{ET-1}}$ for A–C.

that $\text{NEt}_3\cdot 3\text{HF}$ is a corrosive chemical composed of up to 30% hydrofluoric acid. When handling this chemical, butyl rubber gloves should be worn, and an HF spill kit should be kept nearby. The reactions were first performed at a constant potential of 1.1–1.2 V (vs Ag wire) using 10 mol % of A, B, or C (Table 3, entries 1–4). Under these conditions, the unmediated direct electrolysis reaction afforded low (<10%) conversion of **1** and only traces (4%) of product **2**, indicating that direct electrolysis is inefficient at this potential. In contrast, with 10 mol % of A or C under otherwise identical conditions, the yield of **2** was 68 and 83%, respectively, demonstrating effective mediation. A significantly lower yield (15%) was obtained using 10 mol % of B. The moderate performance of B likely reflects a combination of the relatively slow $k_{\text{ET-1}}$ between **1** and B and the modest stability of **1** toward $\text{NEt}_3\cdot 3\text{HF}$ (Table 2, entry 5).

Table 3. A–F-Catalyzed Anodic Fluorination of Dithiane 1 with NEt₃·3HF

entry	conditions	mediator	mol %	yield 3 (%)
1	CPE, 1.2 V	none	--	4
2	CPE, 1.1 V	A	10	68
3	CPE, 1.2 V	B	10	15
4	CPE, 1.2 V	C	10	83
5	CCE, 3 mA	none	--	16
6	CCE, 3 mA	A	10	79
7	CCE, 3 mA	B	10	45
8	CCE, 3 mA	C	10	80
9	CCE, 3 mA	D	10	46
10	CCE, 3 mA	E	10	83
11	CCE, 3 mA	F	10	68
12	CCE, 3 mA	A	1	37
13	CCE, 3 mA	C	1	90
14	CCE, 3 mA	E	1	73



Controlled potential electrolysis (CPE) is limited by relatively low current densities and long reaction times. These are particularly problematic when NEt₃·3HF, which slowly dissolves glass components (e.g., the reference electrode and fritted separator). As such, we also examined the conversion of 1 to 2 using controlled current electrolysis (CCE), with a current density of 3 mA (Table 3, entries 5–8). Under these conditions, the direct electrolysis reaction afforded a low (16%) yield of 2. Significantly higher yields were obtained in the presence of 10 mol % of the mediators, with A (79%) and C (80%) outperforming B (45%).


The modular synthesis of TAC derivatives allows for the facile installation of sterically diverse amino groups on the cyclopropenium core without significant modification of the redox potential.⁵ This enabled the exploration of several additional cyclopropenium mediators (D–F) under analogous conditions. As shown in Table 3, entries 9–11, the structure of the TAC significantly impacts catalytic performance, with yields ranging from 46 to 83%. Using the three top-performing mediators (A, C, and E), we dropped the loading to 1 mol %. The yield of 2 decreased substantially with A and E at this lower loading but remained high (90%) with C. This finding highlights the importance of high k_{ET-1} for C, as it enables faster catalyst turnover and thus lower catalyst loading.

We next turned our focus to the A–C-catalyzed oxidation of 4-methoxybenzyl alcohol 3 to aldehyde 4. Little and co-workers studied electron transfer rates between 3 and mediators A and B in the presence of lutidine, revealing k_{ET-3} values of 6.11 and 74.76 M⁻¹ s⁻¹ for A and B, respectively.^{4c} However, when we attempted an analogous

experiment with C, we observed the complete disappearance of the reversible redox couple. Under these conditions, the anodic scan no longer showed a diffusion-limited current response (Figure S8), implying the decomposition of C. Consistent with these observations, preparative scale C-catalyzed oxidation of 3 under Little's conditions (2 equiv of lutidine in 0.2 M LiClO₄ in 1:4 CH₂Cl₂/MeCN) showed no catalysis. Instead, the observed yield of 4 was within the error of that for direct electrolysis (38% vs 42%, respectively). These results are consistent with the CV studies showing poor stability of C in the presence of lutidine (Table 2, entry 9) and reflect a key limitation of C as a mediator relative to A and B.

We hypothesized that this instability stems from the fast decomposition of radical dication C²⁺ in the presence of lutidine. This was confirmed by subjecting an isolated sample⁸ of C²⁺ (5 mM in CH₃CN) to 1 equiv of lutidine, which resulted in a dissipation of the characteristic dark red color of C²⁺ within 1 min at 25 °C (Figure S11). We reasoned that a less soluble base might exhibit improved compatibility with C²⁺. Indeed, upon screening a variety of inorganic bases, we found that subjecting C²⁺ to 1 equiv of Cs₂CO₃ under otherwise analogous conditions resulted in minimal decay over 2 h at 25 °C (Figure S11).

The use of a heterogeneous base makes it challenging to determine k_{ET-3} using CV. Thus, we moved directly to evaluate preparative-scale oxidations of 3 to 4 with Cs₂CO₃ as the base (Table 4). These reactions were conducted in a divided cell

Table 4. A–C-Catalyzed Benzylic Oxidation of 4-Methoxybenzyl Alcohol

entry	mediator	yield 4 (%)
1	none	29
2	A	70
3	B	1
4	C	70

equipped with reticulated vitreous carbon (RVC) electrodes in 0.1 M KPF₆ in MeCN with 10 mol % of mediators A–C. Upon constant potential electrolysis (1.1 V vs Ag wire) over 5 h, the unmediated reaction passed 2.5 F/mol of charge and afforded a moderate 29% yield of 4. In contrast, with 10 mol % of A or C, the reactions passed a total of 3.7 and 3.6 F/mol, respectively, and both afforded a 70% yield of 4. Interestingly, while B outperforms A and C with lutidine as the base,⁴ under these Cs₂CO₃ conditions, B affords a very low yield (1%), with the corresponding current response never exceeding 1 mA. These results highlight the value of having different mediators with similar potentials, as even relatively small changes to the reaction conditions can have a dramatic impact on mediator performance and compatibility.¹⁵

In conclusion, this report compares trisaminocyclopropeniums to two known redox mediators for several electroorganic transformations. We show that the TACs exhibit electrochemical properties (diffusion rates and electron transfer rates) and catalytic performance similar or superior to triarylamine and triarylimidazole derivatives with comparable redox potentials. We note that the TACs are poorly

compatible with the soluble base lutidine; however, this issue can be mitigated by changing to a less soluble base such as Cs_2CO_3 . We anticipate that these studies will motivate the application of TACs and other readily accessible aminocyclopropenium derivatives to a wider variety of electro-organic transformations.

■ ASSOCIATED CONTENT

Data Availability Statement

The data underlying this study are available in the published article and its [Supporting Information](#).

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.joc.4c00422>.

General procedures for cyclic voltammetry studies and preparative scale electrolyses ([PDF](#))

■ AUTHOR INFORMATION

Corresponding Authors

Long Luo – Department of Chemistry, Wayne State University, Detroit, Michigan 48202, United States; orcid.org/0000-0001-5771-6892; Email: long.luo@wayne.edu

Melanie S. Sanford – Department of Chemistry, University of Michigan, Ann Arbor, Michigan 48109, United States; orcid.org/0000-0001-9342-9436; Email: mssanfor@umich.edu

Authors

Sabrina N. Carneiro – Department of Chemistry, University of Michigan, Ann Arbor, Michigan 48109, United States

Joshua D. Laffoon – Department of Chemistry, University of Michigan, Ann Arbor, Michigan 48109, United States; Present Address: Core R&D Engineering & Process Science, Dow, Midland, Michigan 48667, United States

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.joc.4c00422>

Funding

This work was supported by the NSF Center for Synthetic Organic Electrochemistry, CHE-2002158. JDL was funded by the Arnold O. Beckman Postdoctoral Fellowship in Chemical Sciences.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the NSF Center for Synthetic Organic Electrochemistry, CHE-2002158 (M.S.S., L.L.), and the Arnold and Mabel Beckman Foundation (postdoctoral fellowship to J.D.L.) for their support.

■ REFERENCES

(1) (a) Moeller, K. Synthetic applications of anodic electrochemistry. *Tetrahedron* **2000**, *56*, 9527–9554. (b) Wiebe, A.; Gieshoff, T.; Möhle, S.; Rodrigo, E.; Zirbes, M.; Waldvogel, S. R. Electrifying organic synthesis. *Angew. Chem., Int. Ed.* **2018**, *57*, 5594–5619. (c) Yan, M.; Kawamata, Y.; Baran, P. S. Synthetic organic electrochemical methods since 2000: on the verge of a renaissance. *Chem. Rev.* **2017**, *117* (21), 13230–13319. (d) Kingston, C.; Palkowitz, M. D.; Takahira, Y.; Vantourout, J. C.; Peters, B. K.; Kawamata, Y.; Baran, P. S. A survival guide for the “electro-curious.

Acc. Chem. Res. **2020**, *53*, 72–83. (e) Leech, M. C.; Garcia, A. D.; Pettit, A.; Dobbs, A. P.; Lam, K. Organic electro-synthesis: from academia to industry. *React. Chem. Eng.* **2020**, *5*, 977–990. (f) Hioki, Y.; Costantini, M.; Griffin, J.; Harper, K. C.; Merini, M. P.; Nissl, B.; Kawamata, Y.; Baran, P. S. Overcoming the limitations of Kolbe coupling with waveform-controlled electro-synthesis. *Science* **2023**, *380*, 81–87.

(2) (a) Ogibin, Y. N.; Elinson, M. N.; Nikishin, G. I. Mediator oxidation systems in organic electro-synthesis. *Russ. Chem. Rev.* **2009**, *78*, 89–140. (b) Francke, R.; Little, R. D. Redox catalysis in organic electro-synthesis: basic principles and recent developments. *Chem. Soc. Rev.* **2014**, *43*, 2492. (c) Luca, O. R.; Gustafson, J. L.; Maddox, S. M.; Fenwick, A. Q.; Smith, D. C. Catalysis by electrons and holes: formal potential scales and preparative organic electrochemistry. *Org. Chem. Front.* **2015**, *2*, 823–848. (d) Cardinale, L.; Stahl, S. S.; Kalyani, D.; Lehnher, D. Overview of outer-sphere electron transfer mediators for electro-synthesis. In *Adv. Catal.*; Hevia, E.; Pérez-Temprano, M. H.; Diéguez, M., Eds.; New Horizons in Modern Catalysis: Five Different Perspectives; Academic Press, 2023; Vol. 72, Chapter 2; pp 57–102. (e) Francke, R.; Little, R. D. Redox catalysis in organic electro-synthesis: basic principles and recent developments. *Chem. Soc. Rev.* **2014**, *43*, 2492. (f) Steckhan, E. Indirect electroorganic syntheses – a modern chapter of organic electrochemistry [new synthetic methods (59)]. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 683–701.

(3) (a) Fuchigami, T.; Mitomo, K.; Ishii, H.; Konno, A. Electrolytic partial fluorination of organic compounds: Part 44. Anodic gem-difluorodesulfurization using triarylamine mediators. *J. Electroanal. Chem.* **2001**, *507*, 30–33. (b) Fuchigami, T.; Tetsu, M.; Tajima, T.; Ishii, H. Indirect Anodic Monofluorodesulfurization of β -Phenyl-sulfenyl β -lactams Using a Triarylamine Mediator. *Synlett* **2001**, *2001*, 1269–1271.

(4) (a) Zeng, C.; Zhang, N.; Lam, C. M.; Little, R. D. Novel triarylimidazole redox catalysts: Synthesis, electrochemical properties, and applicability to electrooxidative C–H activation. *Org. Lett.* **2012**, *14*, 1314–1317. (b) Francke, R.; Little, R. D. Optimizing electron transfer mediators based on arylimidazoles by ring fusion: Synthesis, electrochemistry, and computational analysis of 2-aryl-1-methylphenanthro[9,10-d]imidazoles. *J. Am. Chem. Soc.* **2014**, *136*, 427–435. (c) Lu, N.; Yoo, S. J.; Li, L.-J.; Zeng, C.-C.; Little, R. D. A comparative study of organic electron transfer redox mediators: Electron transfer kinetics for triarylimidazole and triarylamine mediators in the oxidation of 4-methoxybenzyl alcohol. *Electrochim. Acta* **2014**, *142*, 254–260.

(5) Walser-Kuntz, R.; Yan, Y.; Sigman, M. S.; Sanford, M. S. A physical organic chemistry approach to developing cyclopropenium-based energy storage materials for redox flow batteries. *Acc. Chem. Res.* **2023**, *56*, 1239–1250.

(6) (a) Sevov, C. S.; Samaroo, S. K.; Sanford, M. S. Cyclopropenium salts as cyclable high potential catholytes in non-aqueous media. *Adv. Energy Mater.* **2017**, *7*, 1602027. (b) Yan, Y.; Robinson, S. G.; Sigman, M. S.; Sanford, M. S. Mechanism-based design of a high-potential catholyte enables a 3.2 V all-organic non-aqueous redox flow battery. *J. Am. Chem. Soc.* **2019**, *141*, 15301–15306. (c) Yan, Y.; Vaid, T. P.; Sanford, M. S. Bis(diisopropylamino)cyclopropenium-arene Cations as High Oxidation Potential and High Stability Catholytes for Non-aqueous Redox Flow Batteries. *J. Am. Chem. Soc.* **2020**, *142*, 17564–17571.

(7) Lambert has shown that trisaminocyclopropenium derivatives are effective mediators in some photoelectrochemical transformations. In these systems, photoexcitation of the radical dication generates a potent photooxidant for various organic transformations. (a) Huang, H.; Strater, Z. M.; Rauch, M.; Shee, J.; Sisto, T. J.; Nuckolls, C.; Lambert, T. H. Electrophotocatalysis with a trisaminocyclopropenium radical dication. *Angew. Chem., Int. Ed.* **2019**, *58*, 13318–13322. (b) Huang, H.; Strater, Z. M.; Lambert, T. H. Electrophotocatalytic C–H functionalization of ethers with high regioselectivity. *J. Am. Chem. Soc.* **2020**, *142*, 1698–1703. (c) Shen, T.; Lambert, T. H. Electrophotocatalytic diamination of vicinal C–H bonds. *Science* **2021**, *371*, 620–626. (d) Shen, T.; Lambert, T. H. C–H amination

via electrophotocatalytic Ritter-type reaction. *J. Am. Chem. Soc.* **2021**, *143*, 8597–8602. (e) Huang, H.; Lambert, T. H. Electrophotocatalytic acetoxyhydroxylation of aryl olefins. *J. Am. Chem. Soc.* **2021**, *143*, 7247–7252.

(8) Robinson, S. G.; Yan, Y.; Hendriks, K. H.; Sanford, M. S.; Sigman, M. S. Developing a Predictive Solubility Model for Monomeric and Oligomeric Cyclopropenium-Based Flow Battery Catholytes. *J. Am. Chem. Soc.* **2019**, *141*, 10171–10176.

(9) (a) Fuchigami, T.; Fujita, T. Electrolytic partial fluorination of organic compounds. 14. The first electrosynthesis of hypervalent iodobenzene difluoride derivatives and its application to indirect anodic gem-difluorination. *J. Org. Chem.* **1994**, *59*, 7190–7192.

(b) Fujita, T.; Fuchigami, T. Electrolytic partial fluorination of organic compounds. 20. Electrosynthesis of novel hypervalent iodobenzene chlorofluoride derivatives and its application to indirect anodic gem-difluorination. *Tetrahedron Lett.* **1996**, *37*, 4725–4728.

(c) Fuchigami, T.; Mitomo, K.; Ishii, H.; Konno, A. Electrolytic partial fluorination of organic compounds: Part 44. Anodic gem-difluorodesulfurization using triarylamine mediators. *J. Electroanal. Chem.* **2001**, *507*, 30–33.

(10) (a) Little and co-workers have demonstrated that the radical cation stability of mediator B is largely dictated by the distortion of the arenes in the triarylimidazole scaffold. By fusing two of the rings on the mediator, radical cation stability is improved significantly. The resulting phenanthroimidazoles have also been explored in the context of photocatalysis. Enders, P.; Majek, M.; Lam, C. M.; Little, R. D.; Francke, R. How to harness electrochemical mediators for photocatalysis – a systemic approach using the phenanthro[9,10-*d*]imidazole framework as a test case. *ChemCatChem* **2022**, *15*, No. e202200830. (b) Nicholson, R. S. Theory and application of cyclic voltammetry for measurement of electrode reaction kinetics. *Anal. Chem.* **1965**, *37*, 1351–1355.

(11) (a) Bard, J.; Faulkner, L. R. *Electrochemical Methods: Fundamentals and Applications*; Harris, D., Swain, E., 10 Robey, C., Aiello, E., Eds.; John Wiley and Sons, Inc.: US, 2001; pp 230–243.

(12) CV cycling studies were conducted under ambient conditions on the benchtop, using a previously dried solution of electrolyte. Stock solutions of electrolyte were dried over activated molecular sieves for at least one day prior to use.

(13) The % loss in peak height was calculated based on the ratio between the peak anodic current response at the tenth cycle and second cycle. This significantly underestimates the decomposition of mediator B because there is a large loss of current between cycles 1 and 2, particularly with $\text{NEt}_3 \cdot 3\text{HF}$, as shown in Figure S7.

(14) Foot-of-the-wave analysis minimizes interfering phenomena, such as side reactions, at high current densities. Strategies for optimized preparative scale electrolysis may then be devised on this basis.

(15) Using controlled current electrolysis (current density = 3 mA) under otherwise identical conditions, the mediated reactions proceeded to form **4** in yields between 45 and 64%. However, direct electrolysis in the absence of mediator afforded comparable yield of 40%, so the mediators offer minimal advantage under these CCE conditions.